

An Interface View of Directed Sandpile Dynamics

Chun-Chung Chen and Marcel den Nijs

Department of Physics, University of Washington, P.O. Box 351560, Seattle, Washington 98195-1560
(March 30, 2001)

We present a directed unloading sand box type avalanche model, driven by slowly lowering the retaining wall at the bottom of the slope. The avalanche propagation in the two dimensional surface is related to the space-time configurations of one dimensional Kardar-Parisi-Zhang (KPZ) type interface growth dynamics. We express the scaling exponents for the avalanche cluster distributions into that framework. The numerical results agree closely with KPZ scaling, but not perfectly.

PACS numbers: 05.40.-a, 05.65.+b, 05.70.Np, 81.05.Rm

Avalanche phenomena are common in nature [1]. They are characterized by fast relaxation dynamics under a slow driving force. Models that describe such dynamics, e.g., so-called sandpile models, have been studied extensively for more than a decade following the work of Bak *et al* [2]. Directed sandpile models are a special subclass in which relaxation follows a directional rule [3], that is, the propagation of active sites occurs only in one direction and never backfires. The central issue in this type of research is whether the dynamics is critical, such that the avalanche distribution functions are scale invariant (power-law decay), and if so, whether these scaling properties are universal in the same sense as equilibrium critical phenomena. Our understanding of these issues is still restricted. There are only a few exactly soluble models, e.g., some deterministic Abelian directed sandpile models [3], but most insight is still limited to numerical simulation data. The evidence for scaling and universality in other types of non-equilibrium dynamics is less ambiguous: In surface growth (another example of intrinsic critical behavior) several universality classes are well established; e.g., Edwards-Wilkinson (EW) [4] and Kardar-Parisi-Zhang (KPZ) [5] type surface growth; population and catalysis type dynamics undergo absorbing state type phase transitions with distinct universality classes like directed percolation and directed Ising [6–8].

Efforts are under way to link avalanche dynamics to these better understood dynamic phenomena. This ranges from mappings to driven interfaces [9–11] and directed percolation [12,13] to using the concepts of renormalization [14,15] and universality classes [16,17,10]. In this letter we introduce a two dimensional (2D) directed avalanche model in which the scaling exponents of the avalanche dynamics are directly related to the exponents of KPZ growth of a 1D interface.

The physical system we have in mind is a sand box with a movable retaining wall to let out sand from the bottom of the slope, see Fig.(1). The retaining wall is lowered very slowly, such that grains tumble out sporadically forming distinct avalanches instead of a continuous flow. We model the sand surface by continuous height variables $h(x, y)$, with respect to a 2D square lattice,

which is rotated over 45° , meaning that in the even (odd) y rows x takes only even (odd) integer values. The retaining wall is placed at the $y = 0$ boundary. The slope is stabilized by the following constraint. The surface particle in column (x, y) is supported by the two columns $(x \pm 1, y - 1)$ just below it, and must be lower than the lowest of the two increased by an amount s_c ,

$$h(x, y) \leq \min[h(x + 1, y - 1), h(x - 1, y - 1)] + s_c. \quad (1)$$

An avalanche is triggered by selecting the highest site, $(x_i, 0)$, at the $y = 0$ wall boundary (it is the i -th avalanche) and reducing its height by a random amount, $0 < \eta_i < s_c$. This represents the lowering of the retaining wall. Next, all sites that violate the stability condition topple according to the rule,

$$h(x, y) \rightarrow \min[h(x + 1, y - 1), h(x - 1, y - 1)] + \eta_i(x, y), \quad (2)$$

where $0 < \eta_i(x, y) \leq s_c$ are uncorrelated random numbers. This toppling continues until the whole system is stable again. Since the toppling of a site in row y can only effect the stability of two sites immediately above it in row $y + 1$, the sites can be updated row by row starting from the $y = 0$ boundary.

This process is idealized compared to a real unloading sand box in the sense that the toppled grains drop out without disturbing the already stabilized lower regions of the surface. It is possible to justify this as the low gravity or strong bond limit where the falling sand does not gain enough momentum to disturb the stabilized surface on its way out. Without this idealization we loose the directed nature of the dynamics and the following surface growth interpretation.

The row-by-row toppling sequence (2) can be reinterpreted as a dynamic rule for a 1D growing interface, in which the y coordinate plays the role of time. Each stable surface configuration represents a space-time configuration of the 1D interface. Imagine creating an initial stable surface configuration, before the retaining wall starts to drop: choose an arbitrary configuration with all $0 < h(x, 0) \leq s_c$ in row $y = 0$. Next, apply Eq.(2) to all sites in the next row, $y = 1$, to create the next slice

of the surface. Repeat this for all higher rows. The configuration in every row is like a 1D interface evolving in time $t = y$. Fig.2 illustrates how this interface propagates during each time step, $y \rightarrow y + 1$. The upper panel shows the first half of the update (the drawn to the dashed line). This is the deterministic part of the propagation (the $\min[\]$ operator) in Eq.(2). The lower panel illustrates the second half of the update, where the heights increase by a random amount $0 < \eta \leq s_c$. The first step removes material, and the second step deposits particles.

This type of interface dynamics almost certainly belongs to the KPZ universality class. Eq.(2) can be rewritten as

$$h(x, t) = \frac{1}{2} [h(x+1, t-1) + h(x-1, t-1)] - \frac{1}{2} |h(x+1, t-1) - h(x-1, t-1)| + \eta(x, t), \quad (3)$$

which is a discrete version of the KPZ Langevin equation [5],

$$\frac{\partial h}{\partial t} = \nabla^2 h - \frac{\lambda}{2} (\nabla h)^2 + \eta, \quad (4)$$

To be absolutely sure, and also to make sure that λ is large enough such that corrections to scaling from the EW point (at $\lambda = 0$) do not interfere, we checked numerically the behavior of the surface width $\Delta(L, t)$, defined as $\Delta^2 \equiv \langle (h - \langle h \rangle)^2 \rangle$. The scaling properties of KPZ growth are known exactly in 1+1 dimensions. Starting from a flat surface at $y = 0$, the width increases as $\Delta \sim t^\beta$ for $0 \ll t \ll L^z$ and saturates at $\Delta \sim L^\alpha$ for $t \gg L^z$, with L the lattice size in the x direction. The exponents are exactly equal to $\alpha = \frac{1}{2}$, $\beta = \frac{1}{3}$ and $z = \alpha/\beta = \frac{3}{2}$. Our numerical results are shown in Fig.(3) and are consistent with the KPZ values.

Throughout this paper we present numerical results in terms of finite size scaling plots of effective exponents, like $\alpha(L)$ in Fig.(3)(a). Global straight line fits to log-log plots of, e.g., $\Delta \sim L^\alpha$ are inaccurate. $\alpha(L)$ is obtained from fitting the form $\Delta \sim L^\alpha$ to two nearby values of L . The approach to $L \rightarrow \infty$ in Fig.(3) is consistent with a leading corrections to scaling exponent $y = -\frac{1}{2}$.

The characteristic feature of self-organized criticality (SOC) is the lack of typical avalanche length, width, depth, or mass scales. The probability distributions follow power-laws, like $P_w \sim w^{-\tau_w}$ characterized by the scaling exponents, τ_l , τ_w , τ_δ and τ_m . Our numerical simulation results confirm the existence of scale invariance. The critical exponents converge well, see Fig.(4). The length l is the maximum y coordinate the avalanche reaches. The width w is the maximum departure in the x direction, $|x - x_i|$, from the trigger point x_i . The depth δ is the maximum height change, $h_i - h_{i-1}$, caused by

the avalanche, and the mass is the total amount of sand carried off by the avalanche.

The meta-distribution function, $P(l, w, \delta)$, should obey the scaling form

$$P(l, w, \delta) = b^{-\sigma} P(b^{-z}l, b^{-1}w, b^{-\alpha}\delta). \quad (5)$$

with b an arbitrary scale parameter. The exponents, σ , z and α , are expected to be robust with respect to details of the dynamic rule, and thus define the universality class. Single parameter distributions, like $P_w \sim w^{-\tau_w}$, follow by integrating out the other two parameters. This implies the following expressions for the τ exponents

$$\tau_l = \frac{\sigma - 1 - \alpha}{z}, \quad \tau_w = \sigma - z - \alpha, \quad \tau_\delta = \frac{\sigma - 1 - z}{\alpha}. \quad (6)$$

or inverted,

$$z = \frac{\tau_w - 1}{\tau_l - 1}, \quad \alpha = \frac{\tau_w - 1}{\tau_\delta - 1}, \quad \sigma = \tau_w + z + \alpha. \quad (7)$$

In Fig.(5) we replot the numerical finite size scaling estimates of the τ exponents in terms of α , z , and σ .

The values $z = 1.52 \pm 0.02$ and $\alpha = 0.46 \pm 0.01$ are very close to those of 1D KPZ growth. Every stable slope configuration represents a possible world sheet of a 1D KPZ type interface, and every avalanche the difference between two such world sheets. Therefore it is natural to expect that the length (depth) of the avalanche scales with the KPZ value for z (α).

The distribution of avalanche cluster sizes is the most commonly studied and experimentally the most accessible property of SOC. Its exponent is linked to α and z in the following manner. At the start of the avalanche, the height of a boundary site ($y = 0$) is lowered on average by $\frac{1}{2}s_c$. Thus, for a sandbox of width, L_x , the boundary row is lowered by $\frac{1}{2}s_c$ after L_x avalanches. In the stationary state, the entire surface moves down on average by $\frac{1}{2}s_c$ and the average amount of removed sand is equal to $L_y \times \frac{1}{2}L_x s_c$. Thus, the average mass of an avalanche must be equal to

$$\langle m \rangle = \int dm m P_m(m) = \frac{1}{2} s_c L_y. \quad (8)$$

This is analogous to conservation of current in conventional deposition type avalanche systems (see, e.g., [3]). Assume that the avalanche clusters are compact, i.e., that the sizes of holes of unaffected regions inside an avalanche do not scale with the avalanche size. In that case, the mass scales as $m \sim l \times w \times \delta$, and we can use the meta-distribution function to evaluate Eq.(8), as

$$\langle m \rangle \sim \int_0^{L_y} dl \int_0^\infty dw \int_0^\infty d\delta l w \delta P(l, w, \delta) + m_{L_y} \int_{L_y}^\infty dl \int_0^\infty dw \int_0^\infty d\delta P(l, w, \delta) \quad (9)$$

This applies when the box is wide and deep enough that L_y is the only limiting factor to the avalanches. The first term accounts for all avalanches that fit inside the box, and the second term for the ones that reach the L_y edge and thus are prematurely terminated. The first integral scales as $L_y^{(-\sigma+2+2z+2\alpha)/z}$ for large L_y , while the second one scales as $L_y^{(-\sigma+1+z+\alpha)/z}$. We have assumed $m \sim l \times w \times \delta$ so $m_{L_y} \sim L_y^{(1+z+\alpha)/z}$. The two terms in Eq.(9) scale in the same way, as

$$\langle m \rangle \sim L_y^{(-\sigma+2+2z+2\alpha)/z}. \quad (10)$$

Solving Eqs.(8) and (10) for σ gives

$$\sigma = 2 + z + 2\alpha \quad (11)$$

Numerically we find $\sigma = 4.43 \pm 0.05$, see Fig.(5), in agreement with the numerical values for α and z , and also with their KPZ values (which yields $\sigma = 9/2$).

In conclusion, we introduced a directed unloading sand box model, in which the stable slope configurations obey KPZ type scaling, and the avalanches represent the difference between two such KPZ world sheets. The avalanche distribution exponents can be reformulated in KPZ language, Eq.(7), and the numerical results agree with the KPZ values.

An intriguing issue for further study is whether the slight systematic deviations from the KPZ values in Fig.(5) are for real or just an artifact of, e.g., finite size scaling (the avalanches cover only a small part of the surface). In KPZ dynamics the scaling exponents follow from an ensemble average over independent MC runs, i.e., over a large set of independent world sheets. The avalanche dynamics performs this ensemble average in a correlated manner. All subsequent world sheets are identical except for the avalanche area. Correlated MC runs like this might well change the scaling exponents (but only slightly apparently), and this might prove a key feature of avalanche type SOC. The dashed curve in Fig.3(b) shows initial results for the global surface roughness in such avalanche correlated MC runs. The surface is noticeably rougher in amplitude but the finite size scaling values of the exponent β are somewhat smaller (consistent with the values for α and z in Fig.5). Whether they will converge in the end to a value $\beta < \frac{1}{3}$ remains to be seen.

In this context it is noteworthy that Eq.(7) applies also to recent solutions to the stochastic directed sandpile model (SDSM) of Paczuski *et.al.* [18] and Kloster *et.al.* [19] where the avalanche dynamics is related to Edwards-Wilkinson (EW) [4] surface growth, with $z = 2$ and $\alpha = 1/2$; and also that the exactly soluble directed sandpile model (DSM) of Dhar and Ramaswamy satisfies Eq.(7) with $z = 2$ and $\alpha = 0$. However, that does not resolve the issue, because both models are intrinsically simpler than ours, and the numerical accuracy of SDSM

in, e.g., Ref. [20,19] is not sufficient to detect a small effect like this.

This research is supported by the National Science Foundation under grant DMR-9985806.

-
- [1] P. Bak, *How nature works: the science of self-organized criticality* (Copernicus, New York, NY, USA, 1996).
 - [2] P. Bak, C. Tang, and K. Wiesenfeld, Phys. Rev. Lett. **59**, 381 (1987).
 - [3] D. Dhar and R. Ramaswamy, Phys. Rev. Lett. **63**, 1659 (1989).
 - [4] S. F. Edwards and D. R. Wilkinson, Proc. R. Soc. London A **381**, 17 (1982).
 - [5] M. Kardar, G. Parisi, and Y.-C. Zhang, Phys. Rev. Lett. **56**, 889 (1986).
 - [6] W. Kinzel, Ann. Isr. Phys. Soc. **5**, 425 (1983).
 - [7] For a review see J. Marro and R. Dickman, *Nonequilibrium Phase Transitions in Lattice Models* (Cambridge University Press, Cambridge, 1996).
 - [8] J. D. Noh, H. Park, and M. den Nijs, Phys. Rev. E **59**, 194 (1999).
 - [9] K. Sneppen, Phys. Rev. Lett. **69**, 3539 (1992).
 - [10] M. Paczuski and S. Boettcher, Phys. Rev. Lett. **77**, 111 (1996).
 - [11] K. B. Lauritsen and M. J. Alava, arXiv.org e-print (1999), cond-mat/9903346.
 - [12] B. Tadić and D. Dhar, Phys. Rev. Lett. **79**, 1519 (1997).
 - [13] A. Vázquez and O. Sotolongo Costa, J. Phys. A **32**, 2633 (1999).
 - [14] L. Pietronero, A. Vespignani, and S. Zapperi, Phys. Rev. Lett. **72**, 1690 (1994).
 - [15] A. Vespignani, S. Zapperi, and L. Pietronero, Phys. Rev. E **51**, 1711 (1995).
 - [16] A. Ben-Hur and O. Biham, Phys. Rev. E **53**, R1317 (1996).
 - [17] A. Mehta, J. M. Luck, and R. J. Needs, Phys. Rev. E **53**, 92 (1996).
 - [18] M. Paczuski and K. E. Bassler, Phys. Rev. E **62**, 5347 (2000).
 - [19] M. Kloster, S. Maslov, and C. Tang, Phys. Rev. E **63**, 026111 (2001).
 - [20] R. Pastor-Satorras and A. Vespignani, Phys. Rev. E **62**, 6195 (2000).




FIG. 1. Sand box with a slowly lowering retaining wall




FIG. 2. KPZ type growth dynamics




FIG. 3. Monte Carlo (MC) results for the global interface width: (a) finite size (L_x) estimates for the saturated surface width exponent α ; (b) finite time estimates for the transient surface width width exponent β from a flat initial configuration. The drawn line represent conventional MC simulations with an uncorrelated world sheet ensemble, and the dashed line are data from avalanche correlated MC runs.




FIG. 4. Finite size scaling estimates of the avalanche distribution exponents, τ_l , τ_w , τ_δ , and τ_m




FIG. 5. Finite size scaling estimates for α , z and σ derived from data in Fig.4 using equation (7).